



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2021

---

## **Efficacy of Retreatment After Failed Direct-acting Antiviral Therapy in Patients With HCV Genotype 1-3 Infections**

Dietz, Julia ; Spengler, Ulrich ; Müllhaupt, Beat ; et al

**Abstract:** Hepatitis C virus infection is causing chronic liver disease, cirrhosis, and hepatocellular carcinoma. By combining direct-acting antivirals (DAAs), high sustained virologic response rates (SVRs) can be achieved. Resistance-associated substitutions (RASs) are commonly observed after DAA failure, and especially nonstructural protein 5A (NS5A) RASs may impact retreatment options.<sup>1-3</sup> Data on retreatment of DAA failure patients using first-generation DAAs are limited.<sup>4-7</sup> Recently, a second-generation protease- and NS5A-inhibitor plus sofosbuvir (voxilaprevir/velpatasvir/sofosbuvir [VOX/VEL/SOF]) was approved for retreatment after DAA failure.<sup>8</sup> However, this and other second-generation regimens are not available in many resource-limited countries or are not reimbursed by regular insurance, and recommendations regarding the selection of retreatment regimens using first-generation DAAs are very important. This study aimed to analyze patients who were re-treated with first-generation DAAs after failure of a DAA combination therapy.

DOI: <https://doi.org/10.1016/j.cgh.2019.10.051>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-183277>

Journal Article

Accepted Version

Originally published at:

Dietz, Julia; Spengler, Ulrich; Müllhaupt, Beat; et al (2021). Efficacy of Retreatment After Failed Direct-acting Antiviral Therapy in Patients With HCV Genotype 1-3 Infections. *Clinical Gastroenterology and Hepatology*, 19(1):195-198.e2.

DOI: <https://doi.org/10.1016/j.cgh.2019.10.051>

# Journal Pre-proof

Efficacy of Retreatment After Failed Direct-acting Antiviral Therapy in Patients With HCV Genotype 1–3 Infections

Julia Dietz, Ulrich Spengler, Beat Müllhaupt, Julian Schulze zur Wiesch, Felix Piecha, Stefan Mauss, Barbara Seegers, Holger Hinrichsen, Christoph Antoni, Perdita Wietzke-Braun, Kai-Henrik Peiffer, Annemarie Berger, Katrin Matschenz, Peter Buggisch, Johanna Backhus, Eugen Zizer, Tobias Boettler, Christoph Neumann-Haefelin, David Semela, Rudolf Stauber, Thomas Berg, Christoph Berg, Stefan Zeuzem, Johannes Vermehren, Christoph Sarrazin, for the European HCV Resistance Study Group



PII: S1542-3565(19)31260-1  
DOI: <https://doi.org/10.1016/j.cgh.2019.10.051>  
Reference: YJCGH 56845

To appear in: *Clinical Gastroenterology and Hepatology*  
Accepted Date: 25 October 2019

Please cite this article as: Dietz J, Spengler U, Müllhaupt B, Schulze zur Wiesch J, Piecha F, Mauss S, Seegers B, Hinrichsen H, Antoni C, Wietzke-Braun P, Peiffer K-H, Berger A, Matschenz K, Buggisch P, Backhus J, Zizer E, Boettler T, Neumann-Haefelin C, Semela D, Stauber R, Berg T, Berg C, Zeuzem S, Vermehren J, Sarrazin C, for the European HCV Resistance Study Group, Efficacy of Retreatment After Failed Direct-acting Antiviral Therapy in Patients With HCV Genotype 1–3 Infections, *Clinical Gastroenterology and Hepatology* (2019), doi: <https://doi.org/10.1016/j.cgh.2019.10.051>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 by the AGA Institute

# **Efficacy of Retreatment After Failed Direct-acting Antiviral Therapy in Patients With HCV Genotype 1–3 Infections**

## **Short title: Retreatment after DAA-failure**

Julia Dietz<sup>1</sup>, Ulrich Spengler<sup>2</sup>, Beat Müllhaupt<sup>3</sup>, Julian Schulze zur Wiesch<sup>4</sup>, Felix Piecha<sup>4</sup>, Stefan Mauss<sup>5</sup>, Barbara Seegers<sup>6</sup>, Holger Hinrichsen<sup>6</sup>, Christoph Antoni<sup>7</sup>, Perdita Wietzke-Braun<sup>8</sup>, Kai-Henrik Peiffer<sup>1</sup>, Annemarie Berger<sup>9</sup>, Katrin Matschenz<sup>10</sup>, Peter Buggisch<sup>10</sup>, Johanna Backhus<sup>11</sup>, Eugen Zizer<sup>11</sup>, Tobias Boettler<sup>12</sup>, Christoph Neumann-Haefelin<sup>12</sup>, David Semela<sup>13</sup>, Rudolf Stauber<sup>14</sup>, Thomas Berg<sup>15</sup>, Christoph Berg<sup>16</sup>, Stefan Zeuzem<sup>1</sup>, Johannes Vermehren<sup>1\*</sup>, Christoph Sarrazin<sup>1, 17\*</sup>, for the European HCV Resistance Study Group

\*these authors contributed equally

<sup>1</sup> Department of Internal Medicine 1, University Hospital Frankfurt, Frankfurt, Germany; German Center for Infection Research (DZIF), External Partner Site Frankfurt, Germany

<sup>2</sup> Department of Internal Medicine I, University of Bonn, Bonn, Germany; German Center for Infection Research (DZIF), Partner-Site Cologne-Bonn, Germany

<sup>3</sup> Swiss Hepato-Pancreato-Biliary Center and Department of Gastroenterology and Hepatology, University Hospital Zürich, Zürich, Switzerland

<sup>4</sup> I. Department of Medicine, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; German Center for Infection Research (DZIF), Partner Site Hamburg-Lübeck-Borstel-Riems, Germany.

<sup>5</sup> Center for HIV and Hepatogastroenterology, Düsseldorf, Germany

<sup>6</sup> Gastroenterologisch-Hepatologisches Zentrum Kiel, Kiel, Germany

<sup>7</sup>Department of Internal Medicine II, University Hospital Mannheim, Mannheim,  
Germany

<sup>8</sup>Medical Department I, University Hospital Schleswig-Holstein, Kiel, Germany

<sup>9</sup>Institute for Medical Virology, University Hospital Frankfurt, Frankfurt, Germany

<sup>10</sup>Institute for Interdisciplinary Medicine IFI, Hamburg, Germany

<sup>11</sup>Department of Internal Medicine I, Ulm University, Ulm, Germany

<sup>12</sup>Department of Medicine II, Medical Center, University of Freiburg, Faculty of  
Medicine, University of Freiburg, Germany

<sup>13</sup>Division of Gastroenterology and Hepatology, Kantonsspital St. Gallen, St. Gallen,  
Schweiz

<sup>14</sup>Department of Internal Medicine, Medical University of Graz, Graz, Austria

<sup>15</sup>Department of Gastroenterology and Rheumatology, University Hospital Leipzig,  
Leipzig, Germany

<sup>16</sup>Department of Internal Medicine I, University of Tübingen, Tübingen, Germany

<sup>17</sup>Medizinische Klinik 2, St. Josefs-Hospital, Wiesbaden, Germany

Corresponding author:

Christoph Sarrazin, MD

Medizinische Klinik 1

Theodor-Stern-Kai 7

D-60590 Frankfurt, Germany

E-Mail: [sarrazin@em.uni-frankfurt.de](mailto:sarrazin@em.uni-frankfurt.de)

Phone: +49-69-6301-87661

Fax: +49-69-6301-84441

**Grant support:** This study was supported by a DZIF (Deutsches Zentrum für Infektionsforschung) grant entitled “HCV Treatment Optimization” to Christoph Sarrazin (TTU 05.809).

**Abbreviations:** DCV, daclatasvir; DAA, direct-acting antiviral; DSV, dasabuvir; EBR, elbasvir; GT, genotype; GLE, glecaprevir; GZR, grazoprevir; ITT, intention-to-treat; LDV, ledipasvir; NS, nonstructural protein; NS3, nonstructural protein 3; NS5A, nonstructural protein 5A, NS5Ai, NS5A inhibitor, NS5B, nonstructural protein 5B, OBV, ombitasvir; PI, protease inhibitor; PIB, pibrentasvir; PTV/R, paritaprevir/ritonavir; PEG-IFN, pegylated interferon; PCR, polymerase chain reaction; RASs, resistance-associated substitutions; SMV, simeprevir, SMV; SOF, sofosbuvir; SVR, sustained virologic response; VEL, velpatasvir; RBV, ribavirin; VOX, voxilaprevir; PrOD (Paritaprevir/Ritonavir/Ombitasvir with Dasabuvir)

**DISCLOSURES:**

**Julia Dietz:** no conflicts to disclose

**Ulrich Spengler:** speaking and/or consulting fees: AbbVie, BMS, Falk, Gilead, Intercept, Merck/MSD.

**Beat Müllhaupt:** Speaking and/or consulting fees: Merck/MSD, AbbVie, Intercept, Astra, Bayer, BMS, Gilead. Research support: Gilead.

**Julian Schulze zur Wiesch:** Speaking and/or consulting fees: AbbVie, Gilead, and Merck/MSD

**Felix Piecha:** no conflicts to disclose

**Stefan Mauss:** Speaking and/or consulting fees: AbbVie, Gilead, Janssen, Merck/MSD

**Barbara Seegers:** no conflicts to disclose

**Holger Hinrichsen:** Speaking and/or consulting fees: AbbVie, BMS, Falk, Gilead, Intercept, Merck/MSD.

**Christoph Antoni:** Speaking and/or consulting fees: AbbVie, Falk, Gilead, Merck/MSD.

**Perdita Wietzke-Braun:** no conflicts to disclose

**Kai-Henrik Peiffer:** no conflicts to disclose

**Annemarie Berger:** no conflicts to disclose

**Katrin Matschenz:** Speaking and/or consulting fees: AbbVie.

**Peter Buggisch:** Speaking and/or consulting fees: AbbVie, BMS, Falk, Gilead, Janssen, Merz Pharma, Merck/MSD.

**Johanna Backhus:** Speaking and/or consulting fees: Gilead, Abbvie, Janssen, Intercept.

**Eugen Zizer:** Speaking and/or consulting fees: AbbVie, Falk, Gilead, Janssen, Intercept

**Tobias Boettler:** Speaking and/or consulting fees: AbbVie, Gilead, Merck/MSD.

**Christoph Neumann-Haefelin:** Speaking and/or consulting fees: AbbVie, Falk Foundation, Gilead.

**David Semela:** Speaking and/or consulting fees: Merck/MSD, Janssen, AbbVie, Boehringer Ingelheim, Intercept, Bayer, BMS, Gilead. Research support: Gilead.

**Rudolf Stauber:** Speaking and/or consulting fees: AbbVie, Bayer, BMS, Gilead, Ipsen, MSD.

**Thomas Berg:** Speaking and/or consulting fees: AbbVie, Alexion, Bayer, Boehringer Ingelheim, BMS, Gilead, GSK, Intercept, Janssen, MSD/Merck, Merz, Novartis, Sequana Medical and Roche. Research support: AbbVie, Roche, BMS, Gilead, Novartis, Merck/MSD, Intercept, Janssen, Novartis, Sequana Medical, and Pfizer.

**Christoph Berg:** Speaking and/or consulting fees: AbbVie, BMS, Gilead, Merck/MSD.

**Stefan Zeuzem:** Speaking and/or consulting fees: Abbvie, BMS, Gilead, Janssen, Merck/MSD.

**Johannes Vermehren:** Speaking and/or consulting fees from Abbott, AbbVie, Gilead, Bristol-Myers Squibb, Medtronic, Merck/MSD, and Roche.

**Christoph Sarrazin:** Speaking and/or consulting fees: Abbott, AbbVie, BMS, Gilead, Intercept, Janssen, Merck/MSD, Roche. Research support: Abbott, Gilead, Janssen, Roche, Siemens. Speaker: Abbott, AbbVie, BMS, Gilead, Intercept, Janssen, Merck/MSD, Roche, Siemens.

**Author contributions:** JD: study concept and design, acquisition of data, analysis and interpretation of data, drafting of the manuscript, critical revision of the manuscript. US: acquisition of data, critical revision of the manuscript. BM: acquisition of data, critical revision of the manuscript. JZW: acquisition of data, critical revision of the manuscript. FP: acquisition of data, critical revision of the manuscript. SM: acquisition of data, critical revision of the manuscript. BS: acquisition of data, critical revision of the manuscript. HH: acquisition of data, critical revision of the manuscript. CA: acquisition of data, critical revision of the manuscript. PWB: acquisition of data, critical revision of the manuscript. KHP: acquisition of data, critical revision of the manuscript. AB: acquisition of data, critical revision of the manuscript. KM: acquisition of data, critical revision of the manuscript. PB: acquisition of data, critical revision of the manuscript. JB: acquisition of data, critical revision of the manuscript. EZ: acquisition of data, critical revision of the manuscript. TB: acquisition of data, critical revision of the manuscript. CNH: acquisition of data, critical revision of the manuscript. DS: acquisition of data, critical revision of the manuscript. RS: acquisition of data, critical revision of the manuscript. TB: acquisition of data, critical revision of the manuscript. CB: acquisition of data, critical revision of the manuscript. SZ: study concept and design, analysis and interpretation of data, critical revision of the manuscript. JV: Study concept and design, acquisition of data, analysis and interpretation of data, drafting of the manuscript, critical revision of the manuscript. CS: Study concept and design, acquisition of data, analysis and interpretation of data, drafting of the manuscript, critical revision of the manuscript, obtained funding.

**Keywords:** Hepatitis C Virus, Resistance-associated substitutions, direct-acting antivirals, retreatment



## INTRODUCTION

Hepatitis C virus (HCV) infection is causing chronic liver disease, cirrhosis and hepatocellular carcinoma (HCC). By combining direct acting antivirals (DAAs), high sustained virologic response rates (SVR) can be achieved. RASs are commonly observed after DAA failure and especially NS5A RASs may impact retreatment options.<sup>1-3</sup> Data on retreatment of DAA failure patients using first generation DAAs are limited.<sup>4-7</sup> Recently, a second generation protease- and NS5A-inhibitor plus sofosbuvir (voxilaprevir/velpatasvir/sofosbuvir, VOX/VEL/SOF) was approved for retreatment after DAA failure.<sup>8</sup> However, this and other second generation regimens are not available in many resource-limited countries or are not reimbursed by regular insurance and recommendations regarding the selection of retreatment regimens using first generation DAAs are very important. This study aimed to analyze patients who were retreated with first generation DAAs after failure to a DAA combination therapy.

## PATIENTS & METHODS

Serum of patients with chronic hepatitis C was collected at different European study sites as part of a resistance database.<sup>2</sup> We searched the database until October 1<sup>st</sup>, 2017 for patients with a DAA failure.

Data was collected retrospectively and recommendations for retreatment were given in line with the EASL/AASLD guidelines. However, individual retreatment strategies were at the treating physician's discretion. A sustained virologic response was defined as negative HCV RNA 12 weeks after end of retreatment (SVR12). We applied a modified intention-to-treat (mITT) analysis including patients with completed retreatment and FU12. The study was conducted in accordance with the

Declaration of Helsinki and approved by the ethics committee of the University Hospital Frankfurt.

HCV NS3, NS5A and NS5B PCR and sequencing analyses were conducted as described previously.<sup>2</sup>

## RESULTS

We investigated a cohort 631 DAA failure patients and 47% (n=262/558) of eligible patients completed retreatment (Supp. Fig. 1).

Overall 84% of NS5Ai/SOF failures infected with GT1 achieved SVR. The majority of patients received a regimen including a PI plus SOF (SMV/SOF±RBV, SMV/SOF+DCV/LDV, PrOD/SOF±RBV, GZR/EBR+SOF) and the SVR rate was 91%. All patients without NS5A RASs achieved SVR, while the SVR rate was 90% in presence of RASs and Y93 RASs reduced SVR rates. A PI-based regimen without SOF (PrOD±RBV, GZR/EBR) was less effective (82% SVR) with slightly lower SVR rates in GT1a compared to 1b. For PI-based regimens in absence of RASs, the SVR rate was 93% and NS5A RASs reduced SVR rates. A repetition of NS5Ai/SOF without a PI was ineffective with 68% SVR. Here, in absence of RAS, the SVR rate reached 88% and was 50% only in presence of RAS (Fig. 1A-D).

All GT3 DCV/SOF failures were retreated NS5Ai/SOF-based and 60% achieved SVR. The majority of patients harbored NS5A RASs (93%), and SVR rates were higher in absence of cirrhosis or absence of Y93H (Fig. 1E/F). We detected no statistically significant differences between SVR and failure patients concerning the presence of cirrhosis and other clinical parameters (Supp. Table 1). Data regarding treatment adherence, liver fibrosis and portal hypertension was not available as

patients with decompensated cirrhosis were not included and therapies were conducted at external centers.

## DISCUSSION

The management of DAA failure patients remains a challenge. Our study comprises a very large European real-world DAA failure cohort and 85% achieved SVR upon retreatment with first generation DAAs.

We demonstrated high SVR rates after NS5Ai/SOF failure in GT1, when SOF was reused in combination with a new DAA class (i.e. switch from a NS5Ai to a PI). However, SVR rates were slightly reduced in presence of high-level resistant Y93 RASs. A retreatment based on the addition of a new DAA class without SOF as well as repetition of NS5Ai/SOF led to reduced SVR rates in presence of NS5A RASs and especially GT1a patients with high-level resistant Q30R responded poorly. These strategies may be considered in the absence of NS5A RASs only.<sup>6</sup>

In DCV/SOF failures with GT3, the prevalence of NS5A RASs as well as cirrhosis was considerably higher compared to GT1. SVR rates to a NS5Ai/SOF repetition, were low especially in presence of cirrhosis or Y93H. Interestingly, in a recent study with inclusion of a PI (VOX/VEL/SOF) the majority of treatment failures had GT3.<sup>8</sup> Thus, retreatment of GT3 remains a challenge.

Limitations of this study include the retrospective data analysis and small patient numbers in specific groups. Taken together, for GT1-infected patients including a new DAA class plus sofosbuvir is highly effective, whereas retreatment of GT3 is requiring multiple targeted regimens including a second generation PI. The results of this real-world study may be of importance for many areas of the world where second generation therapies are not available or not affordable.

## REFERENCES

1. Pawlotsky JM. Hepatitis C Virus Resistance to Direct-Acting Antiviral Drugs in Interferon-Free Regimens. *Gastroenterology*. 2016;151(1):70-86.
2. Dietz J, Susser S, Vermehren J, et al. Patterns of Resistance-Associated Substitutions in Patients With Chronic HCV Infection Following Treatment With Direct-Acting Antivirals. *Gastroenterology*. 2018;154(4):976-88 e4.
3. Di Maio VC, Cento V, Aragri M, et al. Frequent NS5A and multiclass resistance in almost all HCV genotypes at DAA failures: What are the chances for second-line regimens? *Journal of hepatology*. 2018;68(3):597-600.
4. Hezode C, Chevaliez S, Scoazec G, et al. Retreatment with sofosbuvir and simeprevir of patients with hepatitis C virus genotype 1 or 4 who previously failed a daclatasvir-containing regimen. *Hepatology*. 2016;63(6):1809-16.
5. de Ledinghen V, Laforest C, Hezode C, et al. Retreatment With Sofosbuvir Plus Grazoprevir/Elbasvir Plus Ribavirin of Patients With Hepatitis C Virus Genotype 1 or 4 Who Previously Failed an NS5A- or NS3-Containing Regimen: The ANRS HC34 REVENGE Study. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2018;66(7):1013-8.
6. Lawitz E, Flamm S, Yang JC, et al. Retreatment of patients who failed 8 or 12 weeks of Ledipasvir/Sofosbuvir-based regimens with Ledipasvir/Sofosbuvir for 24 weeks. *Journal of hepatology*. 2015;62:S192.
7. Poordad F, Bennett M, Sepe TE, et al. Ombitasvir/paritaprevir/ritonavir and dasabuvir with or without sofosbuvir for patients with hepatitis C virus genotype 1 infection who failed a prior course of direct-acting antiviral therapy. *Journal of Medical Virology*. 2019;91(7):1307-12.
8. Bourliere M, Gordon SC, Flamm SL, et al. Sofosbuvir, Velpatasvir, and Voxilaprevir for Previously Treated HCV Infection. *The New England journal of medicine*. 2017;376(22):2134-46.

## FIGURE LEGENDS:

**Fig. 1: Retreatment in patients with NS5Ai/SOF failure.** SVR rates according to (A) retreatment regimens (GT1); RASs in (B) patients who received a new DAA class plus SOF (GT1); (C) individuals who were retreated with new DAA class without SOF (GT1); (D) patients who received a repetition of NS5Ai/SOF (GT1); (E) retreatment regimens (GT3); RASs in (F) DCV/SOF failures (GT3).

## ACKNOWLEDGEMENTS

We thank Dr. Simone Susser for assistance in the collection and analysis of sequencing and resistance data and Caterina Berkowski, Babithra Yoganathan-Kugarajan, Dany Perner and Sandra Passmann for expert technical assistance.

## Other members of the European HCV Resistance Study group:

**University Hospitals:**; J. Balavoine, E. Giostra (Genf, CH); M. Berning, J. Hampe (Dresden, DE); A. De Gottardi, A. Rauch, N. Semmo (Bern, CH); T. Discher, J. Trauth (Gießen, DE); J. Fischer (Leipzig, DE); M. Gress (Marburg, DE); R. Günther (Kiel, DE); H. Heinzow, J. Schmidt (Münster, DE); A. Herrmann, A. Stallmach (Jena, DE); G. Hilgard, K. Deterding, C. M. Lange, S. Ciesek, H. Wedemeyer (Essen, DE); D. Hoffmann (TU, München, DE); H. Klinker, P. Schulze (Würzburg, DE); F. Kocheise, M. Müller-Schilling (Regensburg, DE); A. Kodal (Lübeck, DE); A. Kremer, M. Ganslmayer, J. Siebler (Erlangen); F. Lammert, J. Rissland (Homburg); M. Löbermann (Rostock, DE); T. Götze, A. Canbay (Magdeburg, DE); A. Lohse, J. von Felden, S. Jordan (Hamburg, DE); A. Maieron (St. Pölten, A); D. Moradpour, J.-P. Chave (Lausanne, CH); C. Moreno (Brüssel, BE); T. Müller, M. Muche, H.-J. Eppe

268 (Berlin); K. Port, T. von Hahn, M. Cornberg, M. Manns (Hannover, DE); L. Reinhardt,  
 269 V. Ellenrieder (Göttingen, DE); J. Rockstroh (Bonn, DE); J. Schattenberg, M. Sprinzl,  
 270 P. Galle (Mainz, DE); E. Roeb (Gießen, DE); M. Steckstor, W. Schmiegeler, N. H.  
 271 Brockmeyer (Bochum, DE); T. Seufferlein (Ulm, DE); W. Stremmel (Heidelberg); B.  
 272 Strey (Duisburg); R. Thimme (Freiburg); A. Teufel, R. Vogelmann, M. Ebert  
 273 (Mannheim, DE); K. Tomasiewicz (Lublin, PL); C. Trautwein, F. Tacke, T. Koenen  
 274 (Aachen, DE); T. Weber (Augsburg); R. Zachoval, J. Mayerle, B. Raziorrouh (LMU,  
 275 München, DE).

276 **Academic Hospitals:** W. Angeli, (Kempten, DE); S. Beckebaum (Essen, DE); C.  
 277 Doberauer (Köln, DE); E. Durmashkina, A. Hackelsberger (Wiesbaden, DE); A.  
 278 Erhardt (Wuppertal, DE); A. Garrido-Lüneburg (Bad Oeynhausen, DE); H. Gattringer  
 279 (Steyr, A); D. Genné (Biel, CH), M. Gschwantler (Vienna, A), F. Gundling (München,  
 280 DE); S. Hametner, R. Schöfl (Linz, A); C. Hartmann (Esslingen, DE); T. Heyer  
 281 (Düren, DE), C. Hirschi (Luzern, CH); A. Jussios (Siegburg); S. Kanzler (Schweinfurt,  
 282 DE); N. Kordecki (Lingen, DE); M. Kraus (Altötting-Burghausen, DE); U. Kullig, S.  
 283 Wollschläger (Dresden, DE); L. Magenta, B. Terziroli Beretta-Piccoli (Locarno, CH);  
 284 M. Menges (Schwäbisch Hall, DE); L. Mohr (Lahr, DE); K. Muehlenberg  
 285 (Regensburg, DE); C. Niederau (Oberhausen, DE); B. Paulweber (Salzburg, A); A.  
 286 Petrides (Bochum, DE); M. Pinkernell (Coesfeld, DE); R. Piso (Olten, CH); W.  
 287 Rambach (Bad Kissingen, DE); M. Reiser (Marl, DE); B. Riecken (Ludwigsburg, DE);  
 288 A. Rieke (Koblenz, DE); J. Roth (Lauterbach, DE); M. Schelling (Lachen, CH); P.  
 289 Schlee (Emden; DE); A. Schneider (Korbach, DE); D. Scholz (Oldenburg, DE); E.  
 290 Schott (Berlin, DE); M. Schuchmann (Konstanz, DE); U. Schulten-Baumer  
 291 (Eggenfelden, DE); A. Seelhoff (Berlin, DE); A. Stich (Würzburg, DE); F. Stickel


292 (Bern, CH); J. Ungemach (Frankfurt, DE); E. Walter (Villingen-Schwenningen, DE); A.  
 293 Weber (Nürnberg, DE); T. Winzer (Bad Hersfeld, DE).  
 294 **Local study sites (private practices), Germany:** W. Abels (Nürnberg); M. Adler  
 295 (Braunschweig); F. Audebert (Regensburg); C. Baermann (Freiburg); E. Bästlein  
 296 (Köln); R. Barth (Nürnberg); K. Barthel (Leipzig); W. Becker (München); J. Behrends  
 297 (Oldenburg); J. Benninger (Regensburg); F. Berger (Düsseldorf); D. Berzow  
 298 (Hamburg); T. Beyer (Rosenheim); M. Bierbaum (Hoyerswerda); O. Blaukat  
 299 (Braunschweig); A. Bodtländer (Frankfurt); G. Böhm (Ludwigshafen); N. Börner  
 300 (Mainz); U. Bohr (Berlin); B. Bokemeyer (Minden); H.R. Bruch (Bonn); D. Bucholz  
 301 (Hagenow); O. Burkhard (Worms); N. Busch (Würselen); C. Chirca (Bad  
 302 Reichenhall); R. Delker (Hamm); J. Diedrich, M. Frank (Marburg); M. Diehl (Hanau);  
 303 A. Dienethal (Frankfurt); P. Dietel (Leipzig); N. Dikopoulos (Dornstadt); M. Dreck  
 304 (Coesfeld); F. Dreher (Rottenburg); L. Drude (Marburg); K. Ende (Erfurt); U. Ehrle, K.  
 305 Baumgartl (Pfungstadt); F. Emke, R. Glosemeyer (Osnabrück); G. Felten, D. Hüppe  
 306 (Herne); J. Fischer (München); U. Fischer (Aalen); D. Frederking (Hamburg); B. Frick  
 307 (Darmstadt); G. Frieze (Gießen); B. Gantke (Düsseldorf); P. Geyer, H. R. Schwind  
 308 (Fulda); M. Glas (Kirchheim unter Teck); T. Glaunsinger (Berlin); F. Goebel  
 309 (München); U. Göbel (Cottbus); B. Görlitz (Augsburg); R. Graf (Leutkirch); H. Gruber  
 310 (Landshut); G. Härter (Ulm); M. Herder (Urbach); T. Heuchel (Chemnitz); S. Heuer  
 311 (Bielefeld); K.-H. Höffl (Traunstein); H. Hörster, J.-U. Sonne (Mönchengladbach); W.  
 312 P. Hofmann (Berlin), F. Holst (Marburg); M. Hunstiger (Augsburg); A. Hurst  
 313 (Darmstadt); E. Jägel-Guedes (München); C. John (Berlin); M. Jung (München); B.  
 314 Kallinowski (Schwetzingen); B. Kapzan (Landshut); W. Kerzel (Forchheim); P.  
 315 Khaykin (Frankfurt); M. Klarhof (Ettlingen); U. Klüppelberg, Wolfratshausen; K.  
 316 Klugewitz (Königs Wusterhausen); B. Knapp (Siegen); U. Knevels (Unna); T.



317 Kochsiek (Schloß Holte-Stukenbrock); A. Körfer (Peine); A. Köster (Friedrichshafen);  
 318 M. Kuhn, A. Langekamp (Kassel); B. Künzig (Waiblingen); R. Link (Offenburg); M.  
 319 Littman (Bad Saulgau); H. Löhr (Wiesbaden); T. Lutz, G. Knecht (Frankfurt); U. Lutz  
 320 (Singen); D. Mainz (Saarlouis); I. Mahle (Freudenstadt); P. Maurer (Bühl); C. Mayer  
 321 (Marburg); V. Meister (Vechta); H. Möller, R. Heyne (Berlin), D. Moritzen (Bielefeld);  
 322 M. Mroß (Berlin); M. Mundlos (Verden); U. Naumann (Berlin); O. Nehls (Stuttgart); K.  
 323 & R. Ningel (Nürtingen); A. Oelmann (Paderborn); H. Olejnik, K. Gadow (Goch); E.  
 324 Pascher (Herrieden); J. Petersen (Hamburg); A. Philipp (Recklinghausen); M. Pichler  
 325 (Stuttgart); F. Polzien (Braunschweig); R. Raddant (Geldern); M. Riedel (Köln); S.  
 326 Rietzler (Albstadt Ebingen); M. Rössle (Freiburg); W. Ruffle (Heidenheim); A. Rump  
 327 (Freiburg); C. Schewe, C. Hoffmann (Hamburg); D. Schleeauf (Berlin); K.J. Schmidt  
 328 (Lübeck); W. Schmidt (Berlin); G. Schmidt-Heinevetter (Bochum); J. Schmidtler-von  
 329 Fabris (Stadtbergen); E. Schnaitmann (Stuttgart), L. Schneider (Fürth); A. Schober,  
 330 S. Niehaus-Hahn (Göttingen); J. Schwenzer (Berlin); T. Seidel (Weimar); G. Seitel  
 331 (Karlsruhe); C. Sick (Bremen); K.G. Simon (Leverkusen); D. Stähler, F. Stenschke  
 332 (Köln); H. Steffens (Berlin); K. Stein (Magdeburg); M. Steinmüller (Ehringshausen); T.  
 333 Sternfeld (Landshut); B. Strey (Duisburg); K. Svensson (Bremen); W. Tacke  
 334 (Königstein); G. Teuber (Frankfurt); K. Teubner (Stuttgart); J. Thieringer (Frankfurt);  
 335 A. Tomesch (Nürnberg); U. Trappe (Hamm); J. Ullrich (Krefeld); G. Urban (Görlitz); S.  
 336 Usadel (Freiburg); A. von Lucadou (Nürnberg); F. Weinberger (Bad Zwischenhahn);  
 337 M. Werheid-Dobers (Bergisch-Gladbach); P. Werner (Böblingen); T. Winter  
 338 (Bielefeld); E. Zehnter (Dortmund); A. Zipf (Mannheim).



Figure 1




**Supporting Table 1.** Characteristics of patients eligible for retreatment infected with HCV genotype 1 (n=251) and genotype 3 (n=63).

	DCV/SOF±RBV, HCV GT1, n=31			LDV/SOF±RBV, HCV GT1, n=220			DCV/SOF, HCV GT3, n=63		
	Re-Tx n=18 (58%)		No Re-Tx <sup>§</sup> n=13 (42%)	Re-Tx n=86 (39%)		No Re-Tx <sup>§</sup> n=134 (61%)	Re-Tx n=15 (24%)		No Re-Tx <sup>§</sup> n=48 (76%)
	SVR n=15 (83%)	FAIL n=3 (17%)		SVR n=72 (84%)	FAIL n=14 (16%)		SVR n=9 (60%)	FAIL n=6 (40%)	
GT1, subtype 1a	7 (47)	2 (67)	9 (69)	36 (50)	7 (50)	69 (52)	-	-	-
GT1, subtype 1b	8 (53)	1 (33)	4 (31)	36 (50)	7 (50)	63 (47)	-	-	-
GT1, subtype other	-	-	-	-	-	2 (1)	-	-	-
GT3, subtype 3a	-	-	-	-	-	-	8 (89)	6 (100)	40 (83)
GT3, subtype 3b	-	-	-	-	-	-	1 (11)	-	1 (2)
GT3, subtype 3h	-	-	-	-	-	-	-	-	1 (2)
GT3, subtype other	-	-	-	-	-	-	-	-	6 (13)
Male gender	13 (87)	2 (67)	8 (62)	56 (78)	12 (86)	98 (73)	8 (89)	6 (100)	41 (85)
Mean Age	53 (33-63)	58 (54-61)	54 (28-71)	59 (36-77)	57 (30-74)	57 (24-84)	51 (37-60)	54 (45-54)	50 (31-65)
Cirrhosis	10 (71) n=14*	2 (67) n=3*	5 (46) n=11*	28 (40) n=70*	8 (57) n=14*	50 (39) n=129*	3 (33) n=9*	5 (83) n=6*	15 (33) n=45*
Prior IFN-experience	9 (69) n=13*	1 (33)	7 (78) n=9*	38 (59) n=64*	6 (55) n=11*	39 (39) n=100*	4 (80) n=5*	5 (100) n=5*	18 (56) n=32*
RBV included in 1 <sup>st</sup> treatment	3 (20)	2 (66)	3 (23)	20 (28)	4 (29)	30 (22)	0 (0)	2 (33)	6 (13)
Duration 1 <sup>st</sup> DAA treatment	n=15*	n=3*	n=11*	n=63*	n=13*	n=107*	n=9*	n=6*	n=40*
8 weeks	-	-	-	20 (32)	3 (23)	37 (35)	-	-	-
12 weeks	7 (47)	2 (67)	7 (64)	36 (57)	10 (77)	61 (57)	7 (78)	4 (67)	28 (70)
24 weeks	8 (53)	1 (33)	3 (27)	7 (11)	-	8 (7)	2 (22)	2 (33)	12 (30)
Early discont.	-	-	1 (9)	-	-	1 (1)	-	-	-
Retreatment			n.a.			n.a.			n.a.
RBV included in Retreatment	13 (87)	1 (33)		38 (53)	9 (64)		5 (71)	5 (83)	
Duration Retreatment									
12 weeks	7 (47)	2 (67)		52 (72)	8 (57)		6 (67)	4 (67)	
24 weeks	8 (53)	1 (33)		19 (26)	5 (38)		3 (33)	2 (33) <sup>+</sup>	
Early discont.	-	-		-	1 (5)		-	-	
Resistance after DAA failure									
Pts. with NS3 RASs <sup>#</sup>	1 (7) n=14	2 (67) n=3*	5 (46) n=11*	14 (20) n=70*	1 (8) n=13*	32 (24) n=133*	- n=5*	- n=5*	- n=18*
Pts. with NS5A RASs <sup>#</sup>	12 (86) n=14*	3 (100) n=3*	5 (46) n=9*	55 (79) n=70*	11 (85) n=13*	107 (84) n=127*	8 (89)	6 (100)	36 (82) n=44*
Pts. with NS3any + NS5Aany	2	1	1	5	-	24	-	-	-
NS3 Q80K	2	1	3	13	-	26	-	-	-
NS3 R155K, A156any, D168any	-	-	2	1	-	6	-	-	-
NS5B S282T	-	-	-	-	-	2	-	-	1

DAA, direct acting antiviral; DCV, daclatasvir; FAIL, failure; IFN, interferon; LDV, ledipasvir; n.a., not applicable; NS3, nonstructural protein 3; NS5A, nonstructural protein 5A; SOF, sofosbuvir; pts., patients; RBV, ribavirin; Re-Tx, retreatment. \*Number of patients with available data; <sup>#</sup>Rate of pts. with RASs; <sup>+</sup>One pt. received 36 weeks LDV/SOF; <sup>§</sup>status of October 2017 (see methods).

Supplementary Figure 1



mITT (modified ITT): Retreatment with completed FU  
 no/non-completed re-tx: with status of October 2017 (see methods).  
 re-tx: retreatment  
 \*patients with FU12 data available  
 \*\*includes n=6 pts. with development of HCC